

Bulletin of the Chicago Academy of Sciences

. . .

A Preliminary Key to the Genera of Clavarioid Fungi

MAXWELL S. DOTY

. - -



. .

Chicago
Published by the Academy
1948

The Bulletin of the Chicago Academy of Sciences was initiated in 1883 and volumes 1 to 4 were published prior to June, 1913. During the following twenty-year period it was not issued. Volumes 1, 2, and 4 contain technical or semi-technical papers on various subjects in the natural sciences. Volume 3 contains museum reports, descriptions of museum exhibits, and announcements.

Publication of the *Bulletin* was resumed in 1934 with volume 5 in the present format. It is now regarded as an outlet for short to moderate-sized original papers on natural history, in its broad sense, by members of the museum staff, members of the Academy, and for papers by other authors which are based in considerable part upon the collections of the Academy. It is edited by the Director of the Museum with the assistance of a committee from the Board of Scientific Governors. The separate numbers are issued at irregular intervals and distributed to libraries and scientific organizations, and to specialists with whom the Academy maintains exchanges. A reserve is set aside for future need as exchanges and the remainder of the edition offered for sale at a nominal price. When a sufficient number of pages have been printed to form a volume of convenient size, a title page, table of contents, and index are supplied to libraries and institutions which receive the entire series.

Howard K. Gloyd, Director of the Museum

Committee on Publications:

Alfred Emerson, Professor of Zoology, University of Chicago.
C. L. Turner, Professor of Zoology, Northwestern University.
Hanford Tiffany, Professor of Botany, Northwestern University.

Bulletin of the Chicago Academy of Sciences

A Preliminary Key to the Genera of Clavarioid Fungi

MAXWELL S. DOTY

The fungi to be treated here are basidiomycetes whose fructifications are erect from their substratum and are, in form, simple clubs or more or less branched structures. The hymenium layer covers all the upper surfaces as a rule; though occasionally it is unilateral or the apex of the fructification may be sterile. The substratum may be soil, humus, wood or undecomposed plant debris. There is evidence that some of the species are mycorrhizal. While the larger clavarioid fungi are almost all fleshy in the sense that a mushroom is of fleshy consistency, some of the smaller species are tough or woody. Though they are all edible if palatable ², usually they are not found in sufficient quantities to make them sought for as food

In pursuing a study of the fungi generally allocated to the genus *Clavaria* and to the family Clavariaceae, the present author has been surprised to find only some half dozen species reported from the state of Illinois. In the neighboring tier of states some three to four dozen species are known. Because of situations such as this no attempt is being made at this time to enumerate *species* but to assist individuals who may collect clavarioid fungi tentative keys to the species known to occur in North America have been prepared. These are under constant revision in typewritten or mimeographed form, and so, while it is hoped they are sometimes helpful to their users, they are not yet worthy of publication. However, to promote study of this group, this paper has been prepared as a key to those *genera* one may expect

2 As far as the author knows there is only one report of poisoning by a member of this group. Roger Heim (11) reported, "*Clavaria formosa* n'est pas comestible; il n'exerce aucune action sur l'estomac, mais agit su l'intestine comme un purgatif."

to find in North America. The numbers in parentheses after each generic name in the key refer to the publications in the list at the end of this paper. These so-designated publications deal with the species of that particular genus in North America, and nearly all the recognized species of all but the more obscure genera, such as *Pistillina*, are covered in these references.

Several of the works cited contain keys to the species but none of them has all the species in the genera as differentiated by the following key. Fries

(1) regular "key" (Clavariaceae) isolated version of the same key as used

fungi imperfecti now excluded. Some workers contemporary to Fries and many workers since have treated his genus, *Clavaria*, as a number of groups that in themselves seem to merit distinction as separate genera. It is hoped that this synoptic artificial key will indicate more or less the relationships among the genera and yet be useful. Some genera of the Hydnaceae (15) and of the Tremellales (14) are of clavarioid form, but while they are basidiomycetes, they are arbitrarily excluded from this treatise.

KEY TO THE NORTH TEMPERATE GENERA OF CLAVARIOID FUNGI

1 Spores hyaline (rarely tinted), mostly smooth and thin-walled; basidia 2- or 4-spored:

2 Fructifications lamellate

Sparassis (17).

2/ Fructifications not lamellate:

3 Spores large (7 microns or over), globose; basidia 2-spored; hymenium white or gray; not staining green with FeSO_4^3 :

4 Fleshy putrescent throughout; no distinct stem portion

Clavulina (5, 6, 8, 17).

4/ Toughish to woody, at least below; with a distinct stem portion LACHNOCLAVULINA section of *Clavulina* (5, 6, 8).

3/ Spores distinctly smaller than above or ellipsoid; basidia typically 4-spored if spores spherical; hymenium often colored otherwise or staining green with FeSO_4 :

5 With stout hyphae in the trama producing setae or gloeocystidia in the hymenium or subhymenium, or the apices of branches truncate or cup-shaped; spores under 8 microns long; not staining green with FeSO_4 :

6/ Apices of some branches truncate to cup-shaped; with gloeocystidia in the hymenium layer

Clavicornia (9).

6 Apices of branches acute; with strong setae in the hymenium or subhymenium

Eriocladius (4).

5/ Without such hyphae or apices; spores various; or staining green with FeSO_4 :

7 Fructifications branched:

8 Finely branched (often under 1 mm.); toughish

Pterula (13).

8/Branches larger; flesh fragile or putrescent:

9 Hymenium green with FeSO_4

Clavariella p. p. (5, 6, 8, 17).

9/ Hymenium not green with FeSO_4

Clavaria p. p. (5, 6, 8, 17).

7/Fructifications simple or rarely branched once above:

10 Fructifications enlarged above or over 1 cm. in diameter above; spores elliptic:

11 Minute (not over 5 mm. tall); or with an abruptly inflated head; not green with FeSO_4 :

12 With an inflated down-turned head

Physalacria (1, 2).

12' With the apex merely enlarged:

13 Hymenium on the expanded blunt apex

Pistillina (17).

13' Hymenium on the sides of the club

Pistillaria p. p. (17, 5).

11/Large (over 2 cm. tall); or turning green with FeSO_4

Clavariadelphus (5, 6, 8, 17).

10/Fructifications not enlarged above (i. e., slenderly clavate to filiform clubs); spores various:

14 Fructifications over 2 cm. tall; not obviously restricted to specific hosts or substrata, or the fructifications fascicled fleshy forms; often with globose spores:

15 Tramal hyphae with many secondary crosswalls; damp connections rare

Clavaria p. p. (5, 6, 8, 17).

15/ Tramal hyphae with secondary crosswalls only rarely; clamp connections on most crosswalls

Clavulinopsis (5, 6, 8, 17).

14/ Fructifications smaller; restricted to specific hosts or substrata which may be sclerotia or living organisms; spores elliptical or flattened on one side:

16 Algal "symbionts," narrowly clavate; mostly on rather bare soil; without a sclerotial base *Gliocoryne* (6, 7, 8).

16/ Not algal "symbionts," filiform or under 2 mm. tall; mostly on dead plant materials:

17 With a sclerotial base; stipe slender and distinct; mostly over 5 mm. tall *Typhula* (18).

17/ Without a sclerotial base; stipe not distinct; mostly less than 2 mm. tall *Pistillaria p. p.* (6, 17).

1/ Spores typically ochraceous, mostly roughened or obdurate-walled ⁴; basidia 4-spored, rarely otherwise:

18 Toughish to woody; spores echinate or sharply warty; hymenium sometimes unilateral; not becoming green with FeSO₄:

19 Coarse, leathery to woody fungi; hymenium often unilateral or branches flattened *Thelephora* (12).

19/ Delicate, toughish to woody fungi; hymenium covering all surfaces of the rounded branches *Scytinopogon* (19).

18/ Fleshy; spores smooth, verrucose to echinulate; hyenium on all lateral surfaces of the branches; becoming green with FeSO₄:

20 Simple unbranched fungi with broadened sterile apices; sometimes "mushroom-like" in form

— *Gomphus p. p.* (3, 6, 8, 16, 17)

20/ branched fungi; "coralloid" in form

Clavariella p. p. (5, 6, 8, 17).

⁴In practice, Clavarioid fungi with spores which under the microscope display walls of appreciable (more than 0.5 microns) thickness, obvious ornamentation or color are placed here.

LITERATURE CITED

1. Baker, Gladys E.
Studies in the genus *Physalacria*. Bull. Torrey Bot. Club, vol. 68, p. 265-268, 105 fig. 1941.
2. Addenda to the genera *Helicogloea* and *Physalacria*. Mycologia, vol. 38, p. 630-638, 25 fig. 1946.
3. Burt, E. A.
The Thelephoraceae of North America. II. *Craterellus*. Missouri Bot. Garden, Annals, vol. 1, p. 327-350, 3 pl. 1914.
4. The Thelephoraceae of North America. XI. *Tulasnella*, *Veluticeps*, *Mycobonia*, *Epithele*, and *Lachnodadium*. Ibid., vol. 6, p. 253-280, 5 pl. 1919.
5. The North American species of *Clavaria*, with illustrations of the type specimens. Ibid., vol. 9, p. 1-78, 11 pl. 1922.

A study of the types of all species recorded from North America with a key for their differentiation. This publication can be obtained from the Missouri Botanical Garden, St. Louis, Mo., for about one dollar.
6. Coker, W. C.
The *Clavarias* of the United States and Canada. 209 pages, 92 plates. University of North Carolina Press, Chapel Hill, N. C. 1923.

The most complete monograph of Clavarioid fungi yet to appear. It has half-tone illustrations of **most** species with line drawings of the spores and hymenial constituents of nearly all the species. The volume may be purchased from the printers for about seven dollars.
7. New or noteworthy Basidiomycetes. Elisha Mitchell Sci. Soc., Journ., vol. 55, p. 373-388, 10 pl. 1939.
8. Doty, Maxwell S.
Clavaria, the species known from Oregon and the Pacific Northwest. Oregon State Monographs, Studies in Botany no. 7, 91 pages, 11 plates. Oregon State College Press, Corvallis, Oregon. 1946.
9. *Clavicornia*, a new genus among the Clavarioid fungi. Lloydia, vol. 10, p. 38-44, 6 fig. 1947.
10. Fries, Elias
Systema Mycologicum I. 520 pages. Lund. 1821.
11. Heim, Roger
Note sur la non comestibilité de *Clavaria formosa* Pers. Soc. Myc. Fr., Bull., vol. 38, p. 175. 1922.
12. Lentz, Paul L.
The genus *Thelephora* in Iowa. Iowa Academy of Science, vol. 49, p. 175-184, 11 fig. 1943.
13. Lloyd, C. G.
The genus *Pterula*. Mycological Writings 5 (note 60), p. 863-870, fig. 1463-1485. 1919.
14. Martin, G. W.
The Tremellales of the north central United States and adjacent Canada. Univ. of Iowa Studies in Nat. Hist., vol. 8, no. 3, p. 1-88, 38 fig. 1944.

15. Miller, L. W., and J. S. Boyle
The Hydnnaceae of Iowa. *Ibid.*, vol. 8, no. 2, p. 1-92, 62 fig. 1943.
16. Murrill, W. A.
Agaricaceae. North American Flora, vol. 9, pt. 3, p. 163-200. 1910.
17. Rea, Carleton
British Basidiomycetae. 799 pages. Cambridge Press, Cambridge, England. 1922.
Probably the last attempt to present the higher basidiomycetous flora for any considerable area in one volume. This is likewise the only extensive publication in English arranged according to the admirable Patouillardian scheme. The work on *Clavaria* closely follows the monograph of that group by Cotton and Wakefield.
18. Remsburg, Ruth E.
Studies in the genus *Typhula*. *Mycologia*, vol. 32, p. 52-96, 58 fig. 1940.
19. Singer, Rolf
New genera of fungi-II. *Lloydia*, vol. 8, p. 139-144. 1945.